

A service of the National Library of Medicine
and the National Institutes of HealthMy NCBI
[Sign In] [Regis]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed for [Go] [Clear]

☒ Limits ☐ Preview/Index ☐ History ☐ Clipboard ☐ Details

Limits: Publication Date to 2002/10/18

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

☐ 1: J Biotechnol. 1994 Feb 14;32(2):107-25.ELSEVIER Links
FULL-TEXT ARTICLE**Evaluation of the proteolytic potential of in vitro-cultivated hybridoma and recombinant mammalian cells.****Kratje RB, Lind W, Wagner R.**

Arbeitsgruppe Zellkulturtechnik, Gesellschaft für Biotechnologische Forschung, Braunschweig, Germany.

The proteolytic potential of culture supernatants derived from recombinant baby hamster kidney (BHK) 21 and mouse-mouse hybridoma cells have been characterized. Several assays using enzyme specific chromogenic artificial peptides, as well as a radioactive test for the detection of the total activity, have been established and were adapted to the special conditions existing in culture media of mammalian cells. Proteolytic activity was detected in human serum albumin containing media which was specific for peptides ending with a terminal arginine. The addition of N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) buffer to the culture media resulted in a significant peptide cleavage potential, supporting the fact that this compound is not recommended as a supplement in animal cell culture media. Medium shock protease activity has been detected in culture supernatants of BHK cells when medium was changed completely, caused by a switch from a serum containing state of growth to a serum-free state of growth which is often used in processes with microcarriers. However, this proteolytic activity showed a transient behaviour whereby its secretion stopped when the cells had adapted to the serum-free medium conditions. Characterization of the proteolytic activities using different specific inhibitors and activators supported the assumption that the proteolytic activity reflects a cell specific composition of proteases which can also change dependent on the culture conditions used.

PMID: 7764558 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort by Send to

Related Links

- Proteolytic activity in the culture supernatants of mouse hybridoma cells. [Dev Biol Stand. 1987]
- Adaptation of BHK-21 cells to growth in shaker culture and subsequent challenge by Japanese encephalitis virus. [Appl Microbiol. 1975]
- Recombinant interferon-gamma secreted by Chinese hamster ovary-320 cells cultivated in suspension in protein-free media is protected against extracellular proteolysis by the expression of natural protease inhibitors and by the addition of plant protein hydrolysates to the culture medium. [In Vitro Cell Dev Biol Anim. 2005]
- Beta-amyloid peptide in regulated secretory vesicles of chromaffin cells: evidence for multiple cysteine proteolytic activities in distinct pathways for beta-secretase activity in chromaffin vesicles. [J Neurochem. 2002]
- Comparison of a production process in a membrane-aerated stirred tank and up to 1000-L airlift bioreactors using BHK-21 cells and chemically defined protein-free medium. [Biotechnol Prog. 2003]

See all Related Articles...

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

☒ Limits ☐ Preview/Index ☐ History ☐ Clipboard ☐ Details

Limits: Publication Date to 2002/10/18

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

☐ 1: Dev Biol Stand. 1987;66:403-8.

Links

Proteolytic activity in the culture supernatants of mouse hybridoma cells.

Schlaeger EJ, Eggimann B, Gast A.

When serum-reduced or serum-free culture supernatants were incubated for 16 hours at 37 degrees C, more than 70-80% of original proteins were digested as measured by gel electrophoretic analysis. The proteolytic activity, which was only observed at pH values lower than 4.5, was reduced in conditioned medium containing higher concentrations of fetal calf serum. During incubation large amounts of the monoclonal antibody (IgG1) were cleaved giving F(ab')₂ fragments. The results reported here strongly suggest that the conditioned medium of mouse hybridoma cultures contain probably two cellular (acid) proteases with apparent MW of 45-50 K and 90-95 K, respectively. The similarities with the lysosomal cathepsin D are discussed.

PMID: 3556271 [PubMed - indexed for MEDLINE]

Related Links

- ▶ Monitoring of the production of monoclonal antibodies by hybridomas. Part II: Characterization and purification of acid proteases present in cell culture supernatants. [Biotechnol. 1991]
- ▶ Evaluation of the proteolytic potential of in vitro-cultivated hybridoma and recombinant mammalian cells. [Biotechnol. 1994]
- ▶ Process-scale purification from cell culture supernatants: monoclonal antibodies. [Dev Biol Stand. 1987]
- ▶ Development of a rapid, single-step procedure using protein G affinity chromatography to deplete fetal calf serum of its IgG and to isolate murine IgG1 monoclonal antibodies from supernatants of hybridoma cells. [Biotechnol. 2000]
- ▶ Antibody production and growth of mouse hybridoma cells in Nutridoma media supplements. [Biotechniques. 1990]

See all Related Articles...

Display AbstractPlus Show 20 Sort by Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

www.pubmed.gov

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

☒ Limits ☐ Preview/Index ☐ History ☐ Clipboard ☐ Details

Limits: Publication Date to 2002/10/18

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

1: J Biotechnol. 1991 Oct;20(3):249-61.

Links

Monitoring of the production of monoclonal antibodies by hybridomas. Part II: Characterization and purification of acid proteases present in cell culture supernatant.

van Erp R, Adorf M, van Sommeren AP, Gribnau TC.

Clinical Lab Systems Research Unit, Organon Teknika B.V., Boxtel, The Netherlands.

An acid proteolytic activity has been found in cell culture supernatants from long-term cultivations of hybridoma cells in hollow fibre bioreactors using serum free medium. The proteolytic activity has now been further characterized and the main results were: (1) the proteolytic activity showed a maximum around pH 3 and declined essentially to zero at pH 8; (2) the activity was specifically inhibited by pepstatin A; (3) the acid proteases consisted of two sets of closely spaced bands with apparent molecular weights of 40-45K and 90-105K, respectively; (4) the protease bands (40-45K and 90-105K) were reactive with anti-human cathepsin D; (5) the IEP values of the acid proteases ranged from pH 4.55-6.5. Furthermore, IgG incubation with the acid proteases isolated from hybridoma cells yielded fragments similar to those found in serum-free hollow fibre cell culture supernatants. These results indicated that the IgG fragments are the result of degradation by cathepsin D like proteases released after cell death or cell lysis.

PMID: 1367572 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort by Send to

Related Links

- ▶ Monitoring of the production of monoclonal antibodies by hybridomas. Part I: Long-term cultivation in hollow fibre bioreactors using serum-free medium. [J Biotechnol. 1991]
- ▶ Proteolytic activity in the culture supernatants of mouse hybridoma cells. [Dev Biol Stand. 1987]
- ▶ Effects of buffering conditions and culture pH on production rates and glycosylation of clinical phase I anti-melanoma mouse IgG3 monoclonal antibody R24. [Biotechnol Bioeng. 2003]
- ▶ Process-scale purification from cell culture supernatants: monoclonal antibodies. [Dev Biol Stand. 1987]
- ▶ Effective production of anti-tetanus toxoid and anti-HBsAg human monoclonal antibodies by serum-free culture of hybridomas. [J Biotechnol. 1991]

See all Related Articles...

Write to the Help Desk
NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer